



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

60. Proposed by Professor C. E. WHITE, Trafalgar, Indiana.

Prove that every algebraic equation can be transformed into another equation of the same degree, but which wants its  $n^{\text{th}}$  term.

61. Proposed by J. A. CALDERHEAD, A. B., Superintendent of Schools, Limaville, Ohio.

Given  $x^2 + x\sqrt{xy} = 10$ , and  $y^2 + y\sqrt{xy} = 20$  to find  $x$  and  $y$  by quadratics.

## CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

### SOLUTIONS OF PROBLEMS.

40. Proposed by F. P. MATZ, D. Sc., Ph.D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

The closed portion of the curve known as "The Cocked Hat," equation

$$x^4 + x^2y^2 + 4ax^2y - 2a^2x^2 + 3a^2y^2 - 4a^3y + a^4 = 0,$$

revolves around the axis of  $y$ . Find the *campanulate* volume generated. If the same portion of the curve revolve around the axis of  $x$ , find the *fusiform* volume generated. Also, determine the area of this closed portion of the curve.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas; W. C. M. BLACK, A. M., Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts; and the PROPOSER.

Solving the equation for  $x^2$  we get  $x^2 = \pm \frac{1}{2}y\sqrt{y^2 + 8ay} - \frac{1}{2}(y^2 + 4ay - 2a^2)$ .

$\therefore$  The campanulate volume generated by the area  $MPA_1QNM$  is


$$V = \frac{\pi}{2} \int_0^a (y\sqrt{y^2 + 8ay} - y^2 - 4ay + 2a^2) dy +$$

$$\frac{\pi}{2} \int_0^a (y\sqrt{y^2 + 8ay} + y^2 + 4ay - 2a^2) dy.$$

$$= \frac{\pi}{2} \left[ \frac{1}{2}(y^2 + 8ay)^{\frac{3}{2}} - 2a(y - 4a)\sqrt{y^2 + 8ay} + 32a^3 \log \right.$$

$$\left. \left\{ y + 4a + \sqrt{y^2 + 8ay} \right\} - \frac{1}{2}y^3 - 2ay^2 + 2a^2y \right]_0^a + \frac{\pi}{2} \left[ \frac{1}{2}y^3 + 2ay^2 - 2a^2y + \frac{1}{2}(y^2 + 8ay)^{\frac{3}{2}} - 2a(y + 4a)\sqrt{y^2 + 8ay} + 32a^3 \log \left\{ y + 4a + \sqrt{y^2 + 8ay} \right\} \right]_0^a$$

$$= \frac{4}{3}\pi a^3(12\log 3 - 13).$$



[ZERR, MATZ, and BLACK.]

From the equation we get  $y = \frac{2a(a^2 - x^2) \pm (a^2 - x^2)\sqrt{a^2 - x^2}}{x^2 + 3a^2}$ .

∴ The fusiform volume is

$$\begin{aligned} V &= 8a\pi \int_0^a \frac{(a^2 - x^2)^{\frac{3}{2}} \sqrt{a^2 - x^2}}{(x^2 + 3a^2)^2} dx = 8\pi a^3 \int_0^{\frac{1}{2}\pi} \frac{\cos^6 \theta d\theta}{(4 - \cos^2 \theta)^2}, \text{ where } x = a \sin \theta, \\ &= 8\pi a^3 \left[ \frac{17\theta}{2} - \frac{44\sqrt{3}}{3} \tan^{-1} \left( \frac{2}{\sqrt{3}} \tan \theta \right) + \frac{\cos \theta \sin \theta}{2} + \frac{8 \sin \theta \cos \theta}{9 \cos^2 \theta + 12 \sin^2 \theta} \right]_0^{\frac{1}{2}\pi} \\ &= 4\pi^2 a^3 \left( \frac{17}{2} - \frac{44\sqrt{3}}{9} \right) = \frac{2}{9} \pi^2 a^3 (153 - 88\sqrt{3}). \end{aligned}$$

$$\begin{aligned} \text{Also area is } A &= 2 \int_0^a \frac{(a^2 - x^2) \sqrt{a^2 - x^2}}{x^2 + 3a^2} dx = 2a^2 \int_0^{\frac{1}{2}\pi} \frac{\cos^4 \theta d\theta}{4 - \cos^2 \theta} = 2a^2 \left[ \frac{8\sqrt{3}}{3} \right. \\ &\left. \tan^{-1} \left( \frac{2}{\sqrt{3}} \tan \theta \right) - \frac{9\theta}{2} - \frac{\sin \theta \cos \theta}{2} \right]_0^{\frac{1}{2}\pi} = \frac{1}{6} \pi a^2 (16\sqrt{3} - 27). \end{aligned}$$

[ZERR, and MATZ.]

$$\begin{aligned} \text{Or, fusiform volume} &= 2\pi \int_0^a (y_1^2 - y_2^2) dx = 16a\pi \int_0^a \frac{(a^2 + x^2)^{\frac{3}{2}} dx}{(x^2 + 3a^2)^2} \\ &= 16a^3 \pi \int_0^{\frac{1}{2}\pi} \frac{\cos^6 \theta d\theta}{(4 - \cos^2 \theta)^2} \\ &= 16a^3 \pi \int_0^{\frac{1}{2}\pi} \left( \frac{\cos^6 \theta}{4} + \frac{2\cos^8 \theta}{4^3} + \frac{3\cos^{10} \theta}{4^4} + \frac{4\cos^{12} \theta}{4^5} + \dots \right) d\theta \\ &= 8\pi^2 a^3 \left\{ \frac{1.3.5}{2.4.6.4^2} + \frac{2.1.3.5.7}{2.4.6.8.4^3} + \frac{3.1.3.5.7.9}{2.4.6.8.10.4^4} + \dots \right\}, \text{ since} \\ \int_0^{\frac{1}{2}\pi} \cos^{2m} x dx &= \frac{1.3.5 \dots (2m-1)}{2.4.6 \dots (2m)} \cdot \frac{\pi}{2}. \end{aligned}$$

$$\text{Area of closed portion} = 2 \int_0^a (y_1 - y_2) dx + 4 \int_0^a \frac{(a^2 - x^2)^{\frac{3}{2}} dx}{x^2 + 3a^2}.$$

$$\begin{aligned} \text{Let } x &= a \sin \theta, A = 4a^2 \int_0^{\frac{1}{2}\pi} \frac{\cos^4 \theta d\theta}{4 - \cos^2 \theta} = 4a^2 \int_0^{\frac{1}{2}\pi} \frac{\cos^4 \theta}{4} + \frac{\cos^6 \theta}{4^2} + \frac{\cos^8 \theta}{4^3} \\ &+ \frac{\cos^{10} \theta}{4^4} + \dots d\theta = 2a^2 \pi \left\{ \frac{1.3}{2.4.4} + \frac{1.3.5}{2.4.6.4^2} + \frac{1.3.5.7}{2.4.6.8.4^3} + \dots \right\}, \end{aligned}$$

which series is also convergent.

[BLACK.]